

 **PERIODICO di MINERALOGIA**  
established in 1930

An International Journal of  
MINERALOGY, CRYSTALLOGRAPHY, GEOCHEMISTRY,  
ORE DEPOSITS, PETROLOGY, VOLCANOLOGY  
and applied topics on Environment, Archaeometry and Cultural Heritage

## Different styles of evolution of CO<sub>2</sub>-rich alkaline magmas: the role of melt composition in carbonate-silicate liquid immiscibility

IRINA P. SOLOVOVA<sup>1\*</sup>, ANDREI V. GIRNIS<sup>1</sup>, IRINA T. RASS<sup>1</sup>, JOERG KELLER<sup>2</sup> and NATALIA N. KONONKOVA<sup>3</sup>

<sup>1</sup> Institute of Geology of Ore Deposits, Petrography, Mineralogy and Geochemistry (IGEM),  
Russian Academy of Sciences, 35 Staromonetny, Moscow, 119017 Russia

<sup>2</sup> Institut für Mineralogie, Petrologie und Geochemie, University of Freiburg, Albertstrasse 23b, 79104 Freiburg, Germany

<sup>3</sup> Institute of Geochemistry and Analytical Chemistry (GEOKHI),  
Russian Academy of Sciences, Kosygina 19, Moscow, 117975 Russia

**ABSTRACT.** — Melt and fluid inclusions were studied in olivine from the olivine melilitite of the Mahlberg complex, Germany. It was shown that the magmas were saturated in CO<sub>2</sub> at a pressure higher than 5 kbar and 1220°C and showed low SiO<sub>2</sub> and high CaO and alkali contents. The magma composition is similar to that of the melilitolites of the Gardiner massif in Greenland (Nielsen *et al.*, 1997), which associate with carbonatite bodies and where evidence for carbonatite-silicate liquid immiscibility was detected in melt inclusions. The absence of carbonatite liquids in Mahlberg is probably related to a relatively high Al<sub>2</sub>O<sub>3</sub> content, which shifted the composition of magma away from the carbonate-silicate liquid immiscibility field.

**RIASSUNTO.** — Inclusioni di fuso e fluide in olivina, sono state studiate in rocce del complesso a olivina melilitite di Mahlberg, Germania. Dallo studio risulta che i magmi erano saturi in CO<sub>2</sub> a pressioni superiori a 5 kb e 1220°C e che contenevano poca SiO<sub>2</sub> ma alti tenori di CaO e alcali. La composizione del magma è simile a quella delle melilitoliti del massiccio di Gardiner in Greenland (Nielsen *et al.*, 1997), a cui sono associati corpi di carbonatiti e vi sono evidenze di immiscibilità allo stato liquido nelle inclusioni di fuso. L'assenza di liquidi carbonatitici a Mahlberg è probabilmente legata al contenuto relativamente alto in Al<sub>2</sub>O<sub>3</sub>, che ha provocato uno spostamento della composizione del magma lontano dal campo di immiscibilità tra liquidi carbonatitici e silicatici.

**KEY WORDS:** *melilitite, Rhinegraben, melt inclusion, carbonatite, liquid immiscibility.*

\* Corresponding author, E-mail: girnis@igem.ru

### INTRODUCTION

Most of igneous carbonatites are thought to result from silicate-carbonate liquid immiscibility during the evolution of alkaline fluid-saturated silicate magma. The role of fluid regime and melt composition is still not fully understood. Because of this, it is instructive to compare carbonatite-bearing and carbonatite-free igneous complexes with similar compositions of silicate rocks. Many igneous carbonatites associate with alkaline rocks dominated by Na or K. There are also examples of carbonatites in association with Ca-rich igneous rocks (melilitites). According to Rass (1998), they were formed from a specific high-Ca initial magma similar to group IB kimberlite (Smith *et al.*, 1985). Nielsen *et al.* (1997) studied melt inclusions in the melilitolites of the carbonatite-bearing Gardiner massif, Greenland and determined the compositions of melts and conditions of carbonatite melt separation. Melilitite-bearing rocks are well known in the Upper Rhinegraben igneous province (Dunworth and Wilson, 1998). This paper presents the results of a study of olivine melilitites from Mahlberg, 100 km north of Keiserstuhl. There is no carbonatite in the Mahlberg massif, which will be compared with carbonatite-bearing melilitite-bearing rocks of the Gardiner.

Our approach is based on the investigation of melt and fluid inclusions in minerals, which bear direct evidence on the composition and conditions of initial and evolved melts. The inclusions were studied using various microscopic heating and cooling stages. Crystal phases and glasses were analyzed by an electron microprobe.

#### PETROGRAPHY AND MINERALOGY

The olivine melilitite sample (XKR 37 OM) studied is composed of olivine and clinopyroxene phenocrysts enclosed by a fine-grained groundmass. Olivine phenocrysts (0.2–1.0 mm, forsterite mole fraction of 0.89–0.92) show wide ranges of CaO (0.18 – 1.1 wt %), NiO (0.05 – 0.32 wt %) and MnO contents (0.15–0.71 wt %), which are linearly dependent of  $mg = Mg/(Mg+Fe)$ . In addition to phenocrysts, there are numerous olivine xenocrysts in the rock. They differ from the phenocrysts in larger size (up to 5 mm), lower  $mg$  (up to 0.81), and lower CaO (0.1–0.2 wt %) and MnO contents (up to 0.3 wt %).

Clinopyroxene occurs as large and small phenocrysts and in the groundmass. The phenocrysts show distinct zoning. Lighter zones in grain margins near late fractures and melt inclusions are enriched in  $Al_2O_3$  (up to 11.7 wt %), FeO (up to 8.7 wt %) and  $TiO_2$  (up to 5.6 wt %) (Table 1). Clinopyroxene from the groundmass show  $mg$  of 0.65. In addition, clinopyroxene occurs as (1) crystalline inclusions in phenocrysts of olivine, (2) crystals in late miarolitic cavities, and (3) daughter minerals in melt inclusions in olivine. Except for  $Na_2O$ , the compositions of all clinopyroxenes show regular changes in element concentrations versus  $mg$  value.

The holocrystalline groundmass consists of olivine, clinopyroxene, melilite, nepheline, phlogopite, Ti-magnetite and perovskite. There are also miarolitic aggregates up to 5 mm across (Fig. 1). They are composed of euhedral clinopyroxene, nepheline, melilite, Ba-rich phlogopite (Dunworth and Wilson, 1998),

perovskite and Ti-magnetite crystals (Table 2) cemented by secondary hydrous minerals (wairakite, lawsonite, and others).

#### INCLUSIONS IN MINERALS

Olivine phenocrysts bear numerous melt inclusions containing a variety of daughter crystals including monticellite (FeO = 10.4 wt%), olivine ( $Fe_{0.88-0.93}$ , up to 1.2 wt % CaO), clinopyroxene, melilite, Ba-rich phlogopite, nepheline, F-bearing apatite and magnetite (with up to 10.5 wt %  $Al_2O_3$ ) (Table 3). The inclusions are partially decrepitated and loosed part of their fluid constituents. Because of this, the inclusions were not homogenized during thermometric experiments. The melting temperatures of daughter phases ranged from 1130–1220°C. Some late and secondary melt inclusions were homogenized at 1160–1100°C.

Clinopyroxene crystals in the miarolitic aggregates contain rare silicate melt and low-density fluid inclusions (Figs. 1c, 1d). The homogenization temperature of the melt inclusions is 1130–1140°C. The density of fluid inclusions is lower than 0.01 g/cm<sup>3</sup>.

Melt in the inclusions are low in  $SiO_2$  and high in CaO (up to 18 wt. %) and alkalis (Table 4). Sodium dominates over potassium in early melts and K/Na ratio increases toward more evolved compositions reaching 1.4. MgO contents are no higher than 9 wt %. The melts are rich in  $P_2O_5$  (up to 1.6 wt %), BaO (up to 0.65 wt %), SrO (up to 0.3 wt %), and F (more than 0.5 wt %). The concentration of Cl was always below the electron microprobe detection limit (0.1 wt %).

In addition to melt inclusions, olivine contains abundant fluid inclusions of the gas ± liquid type (Fig. 2). They are dominated by  $CO_2$  (melting temperature of pure  $CO_2$  = –56.6°C) with a density from 0.30–0.73 g/cm<sup>3</sup>. Similar fluids were detected in gas bubbles of melt inclusions. All fluid inclusions are partially decrepitated.

TABLE 1  
Representative clinopyroxene compositions, wt. %

	1	2	3	4	5	6	7	8
SiO <sub>2</sub>	40.83	47.51	40.52	37.23	44.42	50.66	47.27	43.83
TiO <sub>2</sub>	4.94	2.29	4.69	6.20	3.17	1.96	1.49	4.68
Al <sub>2</sub> O <sub>3</sub>	10.63	5.44	10.68	12.74	7.34	2.60	7.74	8.77
Cr <sub>2</sub> O <sub>3</sub>	0.00	0.15	0.00	0.04	0.00	0.00	0.58	0.67
FeO	8.05	5.73	7.91	9.11	6.94	5.45	4.28	5.20
MnO	0.10	0.09	0.08	0.15	0.10	0.16	0.01	0.05
MgO	10.99	14.50	10.74	9.68	12.85	15.03	15.22	12.47
CaO	24.11	24.38	24.00	23.11	24.34	24.45	21.54	24.10
Na <sub>2</sub> O	0.40	0.27	0.42	0.50	0.34	0.46	1.12	0.71
K <sub>2</sub> O	0.01	0.00	0.00	0.03	0.00	0.00	0.01	0.18
P <sub>2</sub> O <sub>5</sub>	0.02	0.03	0.01	0.03	0.05	0.01	-	-
NiO	0.03	0.03	0.03	0.03	0.00	0.01	-	-
Total	100.10	100.41	99.09	98.84	99.55	100.78	99.26	100.66
mg	0.71	0.82	0.71	0.65	0.77	0.83	0.87	0.81

Note: (1)-(3) Phenocrysts: (1) light green zone, (2) dark green zone, small phenocryst. (4), (5) Groundmass clinopyroxene. (6) Clinopyroxene from miarolitic aggregate. (7) Solid inclusion in olivine. (8) Daughter mineral in melt in olivine.

TABLE 2  
Representative compositions of phases from miarolitic aggregates, wt. %

	Cpx	Ne	Phlg	Ap	Per	Ti-Mt	Mel
SiO <sub>2</sub>	49.56	40.91	29.15	1.36	0.50	2.47	43.08
TiO <sub>2</sub>	1.78	0.04	8.01	0.04	51.26	4.28	0.08
Al <sub>2</sub> O <sub>3</sub>	2.47	34.44	16.29	0.39	0.16	1.42	6.52
FeO	5.41	1.05	7.23	0.40	1.51	82.20	3.27
MgO	14.87	0.18	17.35	0.02	0.16	0.74	8.57
MnO	0.15	0.01	0.12	-	-	0.80	0.07
CaO	23.80	0.56	0.22	52.50	33.26	2.82	33.88
Na <sub>2</sub> O	0.59	14.89	0.22	-	1.84	0.10	3.40
K <sub>2</sub> O	0.19	9.03	4.51	0.02	0.04	0.07	0.18
C <sub>2</sub> O <sub>3</sub>	-	-	-	-	0.05	0.17	-
BaO	-	-	14.98	0.15	-	-	0.07
F	-	-	4.55	8.38	-	-	-
Total	98.82	101.11	102.62	100.95*	88.78**	95.07	99.12

Note: \*Including 37.69 wt% of P<sub>2</sub>O<sub>5</sub>. \*\*Low total of perovskite is due to high REE content (Dunworth, Wilson, 1998).

TABLE 3

*Representative compositions of daughter minerals in melt inclusions, wt.%*

	Mont	OI	Cpx	Ne	Per	Ap	Ti-Mt
SiO <sub>2</sub>	37.05	41.62	41.44	42.20	0.45	1.77	0.90
TiO <sub>2</sub>	0.00	0.00	3.94	0.10	46.78	0.00	13.34
Al <sub>2</sub> O <sub>3</sub>	0.00	0.00	11.95	31.13	0.16	0.00	3.26
FeO	10.41	11.81	8.47	1.99	1.82	0.79	71.72
MgO	20.42	46.94	10.73	1.11	0.18	0.67	6.08
MnO	0.16	0.22	0.05	0.04	0.02	0.00	0.77
CaO	33.23	1.19	24.44	0.96	31.11	52.98	0.06
Na <sub>2</sub> O	0.00	0.00	0.39	13.99	2.75	0.00	0.06
K <sub>2</sub> O	0.00	0.00	0.03	8.94	0.05	0.00	0.12
Cr <sub>2</sub> O <sub>3</sub>	0.11	0.09	0.00	0.00	0.00	0.07	0.58
Total	101.38	101.87	101.44	100.45	83.32	56.28	96.91

TABLE 4

*Representative compositions of melts from inclusions, wt.%*

	1	2	3	4	5	6	7
SiO <sub>2</sub>	38.95	37.89	40.98	37.57	38.99	46.31	69.16
TiO <sub>2</sub>	2.59	3.03	2.05	4.25	2.60	2.07	0.89
Al <sub>2</sub> O <sub>3</sub>	10.22	14.44	15.37	12.55	14.31	18.01	14.12
FeO	10.71	9.18	5.90	10.37	10.44	7.13	2.04
MgO	15.27	8.10	7.76	7.08	7.26	5.80	0.32
MnO	0.19	0.09	0.10	0.13	0.04	0.11	0.03
CaO	14.08	17.74	16.32	19.04	14.61	6.18	0.42
Na <sub>2</sub> O	2.57	6.49	7.36	5.23	5.72	7.18	2.51
K <sub>2</sub> O	1.06	4.23	5.29	3.52	3.57	7.86	5.48
Total	95.64	101.19	101.13	99.74	97.54	100.65	94.97

Note: (1) Rock, (2)-(4) primary inclusions in olivine cores, (5) primary inclusion in olivine rim, (6) secondary melt inclusion in olivine, (7) residual melt in inclusion in Cpx of the miarolitic aggregate.




Fig. 1 – Photomicrographs of mirolitic aggregate in the groundmass of olivine melilitite, transmitted light. (a) General view showing extensive clinopyroxene growth on the walls. The central part is filled with secondary hydrous phases. Scale bar is 1 mm. (b) Fragment of the mirolitic aggregate. Clinopyroxene crystals near the wall. Scale bar is 100  $\mu\text{m}$ . (c) Primary melt inclusion in clinopyroxene with a distinct fluid bubble and a daughter crystal. Scale bar is 10  $\mu\text{m}$ . (d) Primary low-density CO<sub>2</sub> fluid inclusion in clinopyroxene. Scale bar is 10  $\mu\text{m}$ .




Fig. 2 – Partially decrepitated CO<sub>2</sub> inclusions in olivine. (a), (b) Primary liquid (L) + gas (G) CO<sub>2</sub> inclusions surrounded by aureoles of secondary satellite inclusions. CO<sub>2</sub> density is (a) 0.73 g/cm<sup>3</sup>, P = 5 kbar at 1220°C, (b) 0.49 g/cm<sup>3</sup>, P = 1.1 kbar at 1130°C. (c) Pseudo-secondary fluid inclusion (liquid + gas,  $d = 0.33 \text{ g/cm}^3$ , P = 0.7 kbar at 1130°C) with crystals (DM) in liquid CO<sub>2</sub>. (d) Secondary low-density CO<sub>2</sub> fluid inclusion in olivine. Scale bars are 10  $\mu\text{m}$ . L and G are CO<sub>2</sub> liquid and gas, respectively.

## DISCUSSION

The results of melt and fluid inclusions show a considerable role of  $\text{CO}_2$  in the genesis and evolution of the magma. The maximum density of  $\text{CO}_2$  fluid inclusions ( $0.73 \text{ g/cm}^3$ ) suggests a fluid pressure of 5 kbar at  $1220^\circ\text{C}$ . Note that the real pressure was probably even higher, because of the partial loss of fluid components. The distribution of fluid inclusions in phenocrysts suggests at least two stages of phenocryst crystallization differing in the fluid regime. At early stages, cores of olivine and clinopyroxene were formed under relatively stable fluid-saturated conditions. This was followed by a rapid decrease of pressure accompanied by extensive degassing and growth of outer phenocryst zones with abundant fluid inclusions of varying density.

The compositions of the least evolved melts trapped in melt inclusions (Table 4) are characterized by low  $\text{SiO}_2$  and high CaO and alkali contents. CaO-rich melts were reported from melt inclusions in olivine from the olivine-melilite rock of the Guli alkaline carbonatite-bearing complex, Polar Siberia (Rass and Plechov, 2000). Similar melts were found by Nielsen *et al.* (1997) in melilite from Gardiner melilitolite, which shows evidence for silicate-carbonate liquid immiscibility. In contrast, melt inclusions in the Mahlberg melilitite contain no carbonate phases, which is in agreement with the absence of carbonatite occurrence. However, both melt compositions and fluid conditions in Mahlberg were similar to those of Gardiner. There are, however, some important differences in melt compositions at similar MgO contents. The melts from Gardiner are in general lower in  $\text{Al}_2\text{O}_3$  and  $\text{SiO}_2$  (Fig. 3) than those of Mahlberg. Although this difference is not very considerable and comparable with the scatter of microprobe analyses, it could be critical in determining degassing behavior and carbonate stability.

Experimental investigations of silicate-carbonate liquid immiscibility determined a two-liquid field in the  $\text{SiO}_2+\text{Al}_2\text{O}_3+\text{TiO}_2-\text{CaO}+\text{MgO}+\text{FeO}-\text{Na}_2\text{O}+\text{K}_2\text{O}$  ternary

diagram (Fig. 4). On this diagram, the least evolved melts from melilitites of the Gardiner rocks fall within the field of silicate liquid, but their differentiation (melilite fractionation) shifts melt composition toward the two-liquid boundary. This is consistent with the appearance of carbonatite melts at late stages of melt evolution. In contrast, melts from the Mahlberg melilitites were initially richer in Al (and Si) and plot further from the liquid immiscibility field. Moreover, higher Al content at high Na expanded the nepheline stability field and resulted in early crystallization of nepheline (together with olivine) from the melt. The resulting liquid evolution path passes parallel to the liquid immiscibility boundary (fig. 4) and even the most evolved melts remain outside the carbonate-silicate field. Thus, the minute




Fig. 3 – Variations of CaO and  $\text{Al}_2\text{O}_3$  versus  $\text{SiO}_2$  in melts from melt inclusions in minerals of Mahlberg and Gardiner.




Fig. 4 – Different evolution paths of the Gardiner and Mahlberg melts (compositions from melt inclusions) projected into the  $\text{SiO}_2+\text{Al}_2\text{O}_3+\text{TiO}_2 - \text{CaO}+\text{MgO}+\text{FeO} - \text{Na}_2\text{O}+\text{K}_2\text{O}$  plane. The carbonate-silicate liquid immiscibility field is given for  $\text{CO}_2$ -saturated pressures of 2-5 kbar (Freestone, Hamilton, 1980; Kjarsgaard, Peterson, 1991).

differences in melt composition have a dramatic effect on the behavior of volatile components. In the case of Gardiner, the formation of carbonate melts consumed a considerable portion of  $\text{CO}_2$  and the crystallization of evolved melts occurred under subvolcanic conditions. In contrast, the accumulation of  $\text{CO}_2$  in the melt of Mahlberg resulted eventually in rapid degassing and a catastrophic eruption.

#### REFERENCES

- DUNWORTH E.A. and WILSON M. (1998) — *Olivine melilitites of the SW German Tertiary Volcanic Province: Mineralogy and petrogenesis*. *J. Petrol.*, **39**, 1805-1836.
- FREESTONE I.C. and HAMILTON D.L. (1980) — *The*

*role of liquid immiscibility in the genesis of carbonatites – an experimental study*. *Contrib. Mineral. Petrol.*, **3**, 105-117.

- KJARSGAARD B. and PETERSON T. (1991) — *Nephelinite-carbonatite liquid immiscibility at Shombole Volcano, East Africa: petrographic and experimental evidence*. *Mineral. Petrol.*, **43**, 293-314.

NIELSEN T.F.D., SOLOVOVA I.P. and VEKSLER I.V. (1997) — *Parental melts of melilitolites and origin of alkaline carbonatite: evidence from crystallised melt inclusions, Gardiner complex*. *Contrib. Mineral. Petrol.*, **126**, 331-344.

RASS I.T. (1998) — *Geochemical features of carbonatite indicative of the composition, evolution, and differentiation of their mantle magmas*. *Geochem. Int.*, **36**, 137-146.

RASS I.T. and PLECHOV P.Yu. (2000) — *Melt inclusions in olivines from the olivine-melilitite rock of the Guli Massif, northwestern Siberian Platform*. *Dokl. Earth Sci.*, **375**, 389-392.

